

University of Groningen

## De structurele basis van de antilichaamspecifiteit

Seijen, Hendrik Geert

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1967

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Seijen, H. G. (1967). *De structurele basis van de antilichaamspecifiteit*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## SUMMARY

The work described in this thesis was aimed at establishing the chemical basis of antibody specificity, i.e. to answer the question whether antibodies of different specificities possess the same or different amino acid sequences in their combining sites. As elucidation of the total sequences of a number of antibodies seemed beyond reach, we used specific (papain) fragmentation of the antibodies and peptide mapping of the fragments to analyse their structural features.

Chapter I, an introductory chapter, describes relevant features of the antigen-antibody reaction and the structure of immunoglobulin G. Special consideration is given to the different forms of heterogeneity of immunoglobulin G: heterogeneity in electrophoretic and chromatographic properties, in antigenic determinants, and in antigen-binding capacity. The end of the chapter discusses the structural problems of antibodies.

The papain fragments of immunoglobulin G from non-immunized, immune, and hyperimmune rabbits were subjected to fingerprinting after performic acid oxidation and trypsin splitting. The results (Chapter III) clearly show that all antibodies, notwithstanding differences in specificity, to a large extent possess identical amino acid sequences. The number of strongly staining peptides was in agreement with that expected for the four-chain, symmetrical model of immunoglobulin G. The total number of peptides, however, exceeded the expected number; this shows that at least some heterogeneity in amino acid sequence must be present.

The fingerprints of antibodies against a positively charged hapten (p-azo-phenyltrimethylammonium chloride) and against two negatively charged haptens (p-azobenzoic acid and m-azobenzoic acid) (Chapter IV) were, to our surprise, not different from those of normal immunoglobulin G, which is a mixture of a very large number of antibodies. We have thus to conclude that either all antibodies

possess the same amino acid sequences - which we consider, to say the least, highly improbable - or that we failed to detect peptide differences. This failure cannot be due to technical causes, and our haptens were so chosen as to make detectibility of differences maximal. We have, therefore, to conclude (Chapter V) that antibodies against even a single antigenic determinant are populations of molecules with different amino acid sequences in their combining sites.

In our view the antigen acts selectively by inducing the formation of a population of fitting antibodies.